

A topical lotion for the eradication of head lice.

You can use this medicine to treat adults and children over 6 months old without seeing a doctor, but use it carefully to get the best results.

Read all this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist for further advice if live lice are detected after the second treatment with Hedrin®, or if head lice are a persistent or recurring problem.
- If you have any other questions, ask your doctor or pharmacist.
- If you notice any side effects or any side effects not mentioned in this leaflet, talk to your doctor or pharmacist.

In this leaflet

- 1 What is Hedrin® Lotion and what does it do?
- 2 Before using Hedrin®
- 3 How to use Hedrin®
- 4 Possible side effects
- 5 Storing Hedrin®
- 6 Further information

1 What is Hedrin® Lotion and what does it do?

The name of your medicine is Hedrin® 4% Lotion. It is a solution which is highly effective against head lice.

2 Before using Hedrin®

Do not use Hedrin®:

- If you are **allergic** to either of the ingredients listed.

Take special care with Hedrin® Lotion:

- For external use only.
- Avoid contact of the lotion with your eyes. If it accidentally comes into contact with your eyes, flush with water.

Warning:

Keep hair away from sources of ignition, especially naked flames and burning cigarettes whilst being treated with Hedrin®. Treated hair can readily burn if ignited.

If you accidentally swallow some

- If you accidentally swallow some of the solution, you should consult a doctor as soon as possible. Take the pack with you to show which medicine you have swallowed.

Use in pregnancy and breastfeeding:

Hedrin® Lotion may be used during pregnancy and breastfeeding.

Use in infants under 6 months:

Children under 6 months should only be treated under medical supervision.

Driving and using machines:

Hedrin® Lotion does not affect your ability to drive or operate machinery.

3 How to use Hedrin®

Hedrin® is suitable for adults and children aged 6 months and over.

Please note:

If a member of the family has live head lice, check the other family members and treat all other cases of live head lice which are discovered, at the same time.

Hedrin® is for external use only.



Directions

- Apply sufficient lotion evenly over **DRY** hair ensuring that the scalp is fully covered.
- Work into the hair spreading the liquid evenly from the roots to tips. Allow the hair to dry naturally.
- Leave on the hair for at least 8 hours or overnight.
- Wash the hair with normal shampoo, rinse thoroughly with water and dry.

Duration of treatment

- It is important that Hedrin® Lotion is applied again after **seven days** to deal with any lice which may hatch in that time. Failure to repeat the treatment may result in the return of a louse infestation.

4 Possible side effects

Important side effects

If you experience any of the following, stop using Hedrin® immediately and talk to a doctor or pharmacist.

- Allergic reactions such as rash, swelling or weals on the skin.
- Breathing difficulties

Other side effects

When using Hedrin® on broken skin, slight irritation may occur. Sometimes application may cause flaking of the scalp, itching and irritation around the eyes.

If you notice any of these or other side effects, stop using the product and talk to a doctor or pharmacist.

5 Storing Hedrin®

- Keep out of the reach and sight of children.
- This product does not require any other special storage conditions.
- Do not use after the expiry date shown on the pack.
- Care should be taken if Hedrin® is spilled as it may cause a slip hazard.

6 Further information

What Hedrin® Lotion contains

The active substance is Dimeticone 4% w/w. Dimeticone is not a chemical poison. It works by a physical process which effectively covers and drowns the louse.

The other ingredients are Cyclomethicone 5. Cyclomethicone 5 is a carrier ingredient which evaporates after application leaving the active ingredient to kill the lice.

What Hedrin® looks like

It is a clear colourless solution.

Hedrin® is available in 50ml and 100ml packs, although not all packs may be marketed.

Marketing Authorisation Holder and Manufacturer



Thornton & Ross Ltd., Huddersfield, HD7 5QH, UK.

Date of preparation of the package leaflet

This leaflet was prepared in April 2009

Hedrin® is a registered Trade Mark of Thornton & Ross Ltd.

About head lice

- Head lice vary in size according to their age. From the size of a printed full stop.... to the size of a sesame seed  Head lice can be difficult to spot because they can match their colour to that of the environment.
- Lice live close to the scalp where they feed on blood and this can cause the scalp to itch.
- Head lice cannot jump, fly or swim and can only be acquired by direct head to head contact. This is why lice are commonly found on the heads of children and those in close contact with children and why it is important to check the heads of all the family members when head lice are found.
- Head lice only live on humans and have no preference for clean or dirty hair.
- The female louse will lay between 3 and 5 eggs per day and these hatch after 7 days. This is why it is important to repeat this treatment after 7 days to ensure that any lice which have hatched after the first treatment are killed.
- Once the louse has hatched, the empty case (known as a nit) will remain glued to the hair shaft. Nits are completely harmless and can be removed with the fingers or a fine toothed comb.